

ONCODAILY MEDICAL JOURNAL

abstract

Analyzing Patterns of Treatment Gaps and Its Causes to Improve Integrated Care in Pediatric Cancer Patients Undergoing Radiation Therapy at a Tertiary Care University Hospital

Mariam Hina, Maria Tariq, Tooba Ali, Yumna Ahmed, Asim Hafiz, Nasir Ali, Ahmed Nadeem Abbasi, Bilal Mazhar Qureshi

DOI: 10.69690/ODMJ-018-0425-1450



SIOP Asia, 2025, Saudi Arabia

abstract



Analyzing Patterns of Treatment Gaps and Its Causes to Improve Integrated Care in Pediatric Cancer Patients Undergoing Radiation Therapy at a Tertiary Care University Hospital

Authors: Mariam Hina, Maria Tariq, Tooba Ali, Yumna Ahmed, Asim Hafiz, Nasir Ali, Ahmed Nadeem Abbasi, Bilal Mazhar Qureshi

Affiliation: Aga Khan University Hospital (AKUH)
Karachi, Pakistan

DOI: [10.69690/ODMJ-018-0425-1450](https://doi.org/10.69690/ODMJ-018-0425-1450)

Introduction: Objective is to analyze the pattern of treatment gaps and its causes in pediatric cancer patients during radiation therapy (RT) to improve collaborative treatment and streamline the service at a tertiary care university hospital.

Methodology: The hospital database was retrospectively reviewed from January 2022-December 2024 and data was collected for patients <18 years, undergoing RT with or without general anesthesia (GA). Demographic and treatment details were collected along with duration of gap and its underlying cause.

Results: Approximately 177 patients were treated with RT. Mean age was 8.5 ± 4 years with mostly males ($n=112, 63\%$). The common diagnoses were Rhabdomyosarcoma-RMS ($n=36, 20\%$), Ewing Sarcoma-ES ($n=35, 19.77\%$), Hodgkins Lymphoma-HL ($n=30, 17\%$) and Wilms Tumor-WT ($n=26, 15\%$). Relapsed/Refractory HL accounted for ($n=12, 6.78\%$). CNS tumors included Medulloblastoma ($n=3, 2\%$), Germ Cell Tumor-GCT ($n=2, 1\%$) and High-Grade Glioma-HGG ($n=1, 0.56\%$). Forty-five patients (25%) were treated under GA. Most of the patients were treated with

Volumetric Modulated Arc Therapy/VMAT ($n=115, 65\%$). Seventy-nine (45%) patients had no gap during treatment. However, a minimum of 1 day gap ($n=23, 13\%$) and a maximum of 18-day gap was observed ($n=1, 0.5\%$). The mean gap was 3 ± 4 days. Gap was categorized as 1-2 days ($n=35, 20\%$), 3-7 days ($n=41, 23\%$) and >8 days ($n=21, 12\%$). The delay reasons were need for admission and optimization ($n=54, 31\%$), suspicion of disease progression and need for plan re-evaluation ($n=2, 1.12\%$), prolonged chemotherapy at primary center ($n=2, 1.12\%$) and welfare issues ($n=1, 0.5\%$). The most common reason for admission was neutropenic fever ($n=16, 9\%$) followed by chest congestion and transfusion because of decreased cell count, equally ($n=2, 1.1\%$).

Conclusion: Our study shows that almost half of the patients experienced no interruptions, gaps were observed in 55% of cases, with admission due to neutropenic fever being the most common cause. Addressing modifiable factors such as improved pre-treatment optimization and enhancing coordination with referring centers can further streamline RT services and improve outcomes for the patients.