

*abstract*

## **Evaluation of Image-Defined Risk Factors in Neuroblastoma: A 6-Month Retrospective Analysis at a Tertiary Care Hospital**

**Maham Ashraf**

[doi.org/10.69690/ODMJ-018-0425-440](https://doi.org/10.69690/ODMJ-018-0425-440)



**SIOP ASIA 2025  
SAUDI ARABIA**

# ONCODAILY MEDICAL JOURNAL

SIOP ASIA 2025 SAUDI ARABIA

## Evaluation of Image-Defined Risk Factors in Neuroblastoma: A 6-Month Retrospective Analysis at a Tertiary Care Hospital

**Author:** Maham Ashraf<sup>1</sup>

**Affiliation:** <sup>1</sup> University of Child Health Sciences and Children Hospital Lahore

**DOI:** <https://doi.org/10.69690/ODMJ-018-0425-440>

**Introduction:** Neuroblastoma is a common extracranial solid tumor in children. Imaging plays a crucial role in identifying risk factors impacting prognosis, surgical management and treatment. Objective is to evaluate image-defined risk factors in Neuroblastoma patients presenting to our tertiary care hospital over 6 months.

**Methodology:** Retrospective review of imaging studies (PAN CT) of neuroblastoma patients (ages 0-18) from January-June 2024. Image-defined risk factors assessed: tumor size, location, invasion into surrounding structures, metastatic disease, intraspinal extension and encasement of vessels.

**Results:** 25 patients (ages 0-18) with neuroblastoma

- IDRFs present in 15 (60%) patients
- Common IDRFs: spinal canal invasion (40%), vascular encasement (32%), neck and mediastinum involvement (28%)
- Metastatic disease: bone (56%), bone marrow (40%), liver (20%)
- Correlation between IDRFs and poor prognosis ( $p < 0.05$ )

**Conclusion:** Image-defined risk factors predict poor prognosis in neuroblastoma. Radiologic assessment of IDRFs is essential for risk stratification and treatment planning.