

# ONCODAILY MEDICAL JOURNAL

*abstract*

## **Pelvic Resections in Pediatric Patients with Ewing Sarcoma**

**Nikolay Bolshakov, Artem Artemov, Anna Bykova,  
Alexey Vorochai, Elena Konopleva, Nikolay Grachev**

DOI: 10.69690/ODMJ-018-0425-1015



SIOP Asia, 2025, Saudi Arabia

*abstract*



## **Pelvic Resections in Pediatric Patients with Ewing Sarcoma**

**Authors:** Nikolay Bolshakov, Artem Artemov, Anna Bykova, Alexey Vorochai, Elena Konopleva, Nikolay Grachev

**Affiliation:** Dmitry Rogachev National Medical Research Center Of Pediatric Hematology, Oncology and Immunology

**DOI:** 10.69690/ODMJ-018-0425-1015

**Introduction:** To date, the world literature describes very little data on the effectiveness of surgical treatment as part of complex therapy for Ewing sarcomas of the pelvic in the pediatric population.

**Methodology:** 40 patients diagnosed with Ewing sarcoma of the pelvic aged under 18 years were operated on in the period from 2012 to 2024. The mean age was 12,5 years (2,3-17,4 years), follow-up was 4,1 years (0,5-11,9 years). Distant metastases were detected in 46.1%. Lesions of more than 1 segment of the pelvic or tumor length more than 8 cm were diagnosed in 82.0%. All patients received surgical treatment according to treatment protocols (Euro Ewing 2008 or Ewing 2012), as well as local radiation therapy in the postoperative period.

**Results:** Combined resection with removal of more than one segment of the pelvic according to the Enneking classification was performed in 77% of patients. The margins were clear in 94.9% of patients. Reconstruction using a synthetic mesh was performed in 10 patients, a screw-rod reconstruction with bone cement – in 12, a custom 3D endoprosthesis in 5, a modular endoprosthesis (“ice cream spoon” type) in 2 patients.

No reconstruction was performed in 11 patients. The average functional result according to the MSTs scale was 73% (58-87%).

Event-free survival (EFS) for patients with the non-metastatic disease form was  $68 \pm 12\%$ , with the metastatic form –  $71 \pm 11\%$ , overall survival (OS) was  $76 \pm 13$  and  $92 \pm 8$ , respectively. When assessing the EFS and OS depending on the initial metastases ( $p=0.865$ ;  $p=0.439$ , respectively) and tumor localization – above and below the midline of the acetabulum ( $p=0.459$ ;  $p=0.271$ , respectively), no statistically significant difference was found.

**Conclusion:** This work analyzes the use of surgical treatment in pediatric and adolescent patients, demonstrates the possibility of achieving good oncological and functional results.