

abstract

Results of Complex Treatment Rhabdomyosarcoma in Children

**Ezozakhon Azimova, Djamilya Polatova, Tojiddin
Mustafojev, Nargiza Karimova, Farkhod Khaitov, Anvar
Shukrullayev, Husen Saitov**

doi.org/10.69690/ODMJ-018-0425-1317



**SIOP ASIA 2025
SAUDI ARABIA**

ONCODAILY MEDICAL JOURNAL

SIOP ASIA 2025 SAUDI ARABIA

Results of Complex Treatment Rhabdomyosarcoma in Children

Author: Ezozakhon Azimova ¹, Djamilya Polatova, Tojiddin Mustafojev, Nargiza Karimova, Farkhod Khaitov, Anvar Shukrullayev, Husen Saitov

Affiliation: ¹ Scientific-Practical Center of Pediatric Hematology, Oncology and Immunology//Tashkent State Dental Institute

DOI: <https://doi.org/10.69690/ODMJ-018-0425-1317>

Introduction: The portion of rhabdomyosarcomas in pediatric oncopathology is on average 1-2%, among soft tissue sarcomas – 5-6%. Aggressive course, frequent metastasis and recurrence of the disease require a complex approach to treatment. The purpose was to study the effectiveness of complex treatment of rhabdomyosarcomas in children at the Scientific and Practical Medical Center of Pediatric Hematology, Oncology and Immunology, Ministry of Health of the Republic of Uzbekistan.

Methodology: The study included 40 patients with rhabdomyosarcoma. The tumor was localized in the corpus in 15 patients (37.5%), in the extremities in 14 (35%), in the perineum in 3 (7.5%), in the head and neck in 3 (7.5%), in the bladder in 3 (7.5%), and in the pelvis in 2 (5%). The average age of the patients was 4.7 years. All patients were diagnosed with stages II-III of the disease. At the first step, all patients after histological verification, underwent neoadjuvant polychemotherapy from 2 courses to 4 courses (PCT) according to the VAIA III CWS 2012 (I2VAd and I2VA schemes): Vincristine - 1.5 mg / m² intravenously on days 1, 8 and 15; lphosphamide - 3000 mg / m² intravenously on days 1 and 2; Doxorubicin – 2x20 mg / m² intravenously for 1 and 2 days; Actinomicyn D – 1,5 mg / m² intravenously with hyperhydration on the 6th day.

Complete tumor regression was observed in 3 (7,5%), partial regression - in 26 (65%) and in 11(27,5%) without effect. Polychemotherapy courses were carried out with concomitant therapy. At the second step, surgical interventions and for patients over 5 years old additionally radiation therapy were performed. In the postoperative period, up to 6 courses of adjuvant polychemotherapy were carried out. Three-year relapse-free and metastasis-free survival was 64.8%.

ONCODAILY MEDICAL JOURNAL

SIOP ASIA 2025 SAUDI ARABIA

Conclusion: A complex approach to the treatment of rhabdomyosarcomas in children significantly improves outcomes and increases relapse-free and metastasis-free survival of patients.