

# ONCODAILY MEDICAL JOURNAL

*abstract*

## **Periodically intensification during maintenance chemotherapy is effective in pediatric acute lymphoblastic leukemia**

**Fenfen Jin, Xiaojun Xu**

[doi.org/10.69690/ODMJ-018-0425-2205](https://doi.org/10.69690/ODMJ-018-0425-2205)



**SIOP ASIA 2025  
SAUDI ARABIA**

# ONCODAILY MEDICAL JOURNAL

SIOP ASIA 2025 SAUDI ARABIA

**Periodically intensification during maintenance chemotherapy is effective in pediatric acute lymphoblastic leukemia**

**Author:** Fenfen Jin <sup>1</sup>, Xiaojun Xu

**Affiliation:** <sup>1</sup> Children's Hospital Zhejiang University School of Medicine

**DOI:** <https://doi.org/10.69690/ODMJ-018-0425-2205>

**Introduction:** Objective is to analyze the outcome of children with acute lymphoblastic leukemia (ALL) treated with the ZJCH-ALL-2019 protocol, which periodically intensifies during maintenance chemotherapy.

**Methodology:** Retrospective analysis of 751 children treated at the Department of Hematology-Oncology, the Children's Hospital, Zhejiang University School of Medicine from October 2018 to May 2023. Kaplan-Meier and Log Rank tests assessed event-free survival (EFS), relapse-free survival (RFS), and overall survival (OS). Cox regression identified prognostic factors.

**Results:** During this period, 446 males and 305 females were enrolled, with median age of 4.9 years, including 76 T-ALL and 675 B-ALL cases. The overall 5-year EFS rate for this treatment regimen was  $91.8 \pm 1.0\%$ , and the 5-year OS rate was  $95.8 \pm 0.7\%$ . Five-year EFS rates for specific genetic fusions varied, with ETV6-RUNX1 at  $98.3 \pm 1.0\%$ , TCF3-PBX1 at  $83.6 \pm 6.3\%$ , BCR-ABL1 at  $82.2 \pm 10.7\%$ , SIL-TAL1 at  $80.0 \pm 10.3\%$ , and MLLr at  $93.3 \pm 6.4\%$ . Additionally, 202 patients (26.89%) exhibited high hyperdiploidy, correlating with a 5-year EFS of  $95.9 \pm 1.7\%$ . Following risk stratification, the patient cohort was categorized as follows: 290 low-risk patients (38.61%) exhibited a 5-year EFS rate of  $98.8 \pm 0.7\%$ ; 321 intermediate-risk patients (42.74%) had a 5-year EFS rate of  $92.2 \pm 1.8\%$ ; and 140 high-risk patients (18.64%) presented with a 5-year EFS rate of  $82.9 \pm 3.4\%$ . There was a total of 24 deaths without relapse, accounting for 3.1% of the patients, and 30 relapses, representing 3.9% of the cases. The Cox multivariate regression analysis showed that the white blood cell count at initial diagnosis, extramedullary lesions, minimal residual disease (MRD) level on day 15 of induction therapy, fusion genes, and immunophenotyping all had significant impacts on prognosis, with extramedullary lesions and the MRD level on day 15 of induction therapy being the most significant factors.

# ONCODAILY MEDICAL JOURNAL

SIOP ASIA 2025 SAUDI ARABIA

**Conclusion:** Periodically intensification during maintenance chemotherapy has shown remarkable effectiveness, with excellent outcome for pediatric B-ALL.