ONCODAILY MEDICAL JOURNAL

abstract

Treatment outcomes of pineal region lesions: 23-year experience from a tertiary care center

Komal Naeem, Zunaira Saeed, Syed Murtaza Nawab, Afia Salman, Malaika Siddiqui, Shilpa Golani, Haseeb Qadri, Hamza Khan, Syed Ather Enam

DOI: 10.69690/ODMJ-018-0915-5252



ONCODAILY MEDICAL JOURNAL

abstract



Treatment outcomes of pineal region lesions: 23-year experience from a tertiary care center

Authors: Komal Naeem¹, Zunaira Saeed², Syed Murtaza Nawab³, Afia Salman², Malaika Siddiqui⁴, Shilpa Golani³, Haseeb Qadri⁵, Hamza Khan⁶, Syed Ather Enam¹

Affiliation: ¹ Centre of Oncological Research in Surgery, Aga Khan University, Karachi, Pakistan

- ² Department of Surgery, Dow University of Health Sciences, Karachi, Pakistan
- ³ Aga Khan University, Karachi, Pakistan
- ⁴ Section of Neurosurgery, Department of Surgery, Aga Khan University, Karachi, Pakistan
- Department of Neurosurgery, Punjab Institute of Neurosciences, Lahore, Pakistan
- ⁶ Rawalpindi Medical University, Rawalpindi, Pakistan

DOI: 10.69690/ODMJ-018-0915-5252

Introduction: Tumors in the pineal region are exceptionally rare and comprise only 0.5-1.1% of all brain tumors. Pineal region was once thought of as the 'forbidden zone' for surgery due to the significant surgical challenges that it posed, especially due to its deep-seated location. In this study we aim to describe experience from a single tertiary care center of managing pineal region lesions. We analyzed the treatment outcomes of the patient.

Methodology: This is a single-center cross-sectional study. Data for all patients who underwent treatment of pineal region lesions from 2000 to 2023 were included. Relevant data points were extracted from the hospital's medical records. Descriptive analysis of the data was performed.

Results: We enrolled 51 patients with a mean age of 23.18 ± 15.7 years, of them, 40% were ≤ 18 years of age.

There were 26 (51%) were male. Most common presenting complaints included headache (94.1%), nausea/vomiting (49%), and diplopia (19.6%). Preoperative hydrocephalus was reported in 34 (67%) patients. Mean tumor size (largest dimension) was 3.6±1.4 cm. All but 3 underwent surgical resection, most of the patients had two surgical procedures done i.e. biopsy and tumor resection. On average, surgery duration was 266.95±217.63 minutes and mean estimated blood loss was 585.79±603.33 ml. Most common histopathological diagnosis were Glioma (30%), Pineoblastoma (11%), and pineal parenchymal tumor of intermediate differentiation (9%). Patients who underwent surgical resection, 61% had gross or near total resection. An average clinical follow-up of 44.2±56.4 months was calculated.

Conclusion: With technological advances the surgical management of pineal tumors is improving. We reported an average follow-up of 44.2±56.4 months. Patients who underwent surgical resection, 61% had gross or near-total resection. Both supraand infra-tentorial approaches were used depending on the tumor location.

Conflict of Interest: None

Funding: None

Disclosure statement: The authors thank Dr. Saad Akhtar Khan, Ceemal Khan, and Roua Nasir from Neurosurgery Research Alliance for their administrative support.

License: This article is published under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0).

© Komal Naeem, 2025. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.